

DOI: 10.3969/j.issn.1000-3142.2014.03.001

王文采. 东川铁线莲, 云南毛茛科一新种[J]. 广西植物, 2014, 34(3):287—289

WANG Wen-Tsai. *Clematis dongchuanensis*, a new species of Ranunculaceae from Yunnan[J]. Guihaia, 2014, 34(3):287—289

# 东川铁线莲, 云南毛茛科一新种

王文采

(中国科学院植物研究所 系统与进化植物学国家重点实验室, 北京 100093)

**摘要:** 描述了自云南东北部发现的毛茛科铁线莲属一新种, 东川铁线莲。此种由于其叶为单叶, 萼片直立, 雄蕊的花丝和花药均被短柔毛而与曲柄铁线莲近缘, 与后者的区别在于其叶呈心状卵形, 叶缘具粗牙齿, 花序为二歧聚伞花序, 以及花药顶端钝, 不具短尖头。

**关键词:** 毛茛科; 铁线莲属; 新种; 云南**中图分类号:** Q949.746   **文献标识码:** A   **文章编号:** 1000-3142(2014)03-0287-03

## *Clematis dongchuanensis*, a new species of Ranunculaceae from Yunnan

WANG Wen-Tsai

(State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany,

the Chinese Academy of Sciences, Beijing 100093, China)

**Abstract:** A species of the genus *Clematis* (Ranunculaceae), *C. dongchuanensis*, is described as new from northeastern Yunnan. In having simple leaves, erect sepals and puberulous filaments and anthers of stamens this species is related to *C. repens* Finet & Gagnep., and from the latter differs in its cordate-ovate and coarsely dentate leaves, dichotomous cymes, and obtuse, not apiculate apexes of anthers.

**Key words:** Ranunculaceae; *Clematis*; new species; Yunnan

### 东川铁线莲 图 1

#### *Clematis dongchuanensis* W. T. Wang, sp. nov.

Fig. 1.

sect. *Viorna* (Reichb.) Prantl subsect. *Connatae* Koehne

Ob folia simplicia, sepala erecta, et staminum filamenta antherasque omnino puberulas species nova haec est affinis *C. repenti* Finet & Gagnep., quae foliis simplicibus vel folii ternati foliolis ellipticis indivisis basi late cuneatis vel rotundatis raro subcordatis margine sparse denticulatis vel integris, cymis axillaribus 1-floris, sepalis dorso glabris, antheris apice apiculatis facile differt.

Woody vine. Stem 1—2 m long (according to the collector); branchlets slender, ca. 1 mm across, sparsely puberulous. Leaves opposite, simple; blades papery, cordate-ovate or pentagonal-ovate, 9—12 cm long, 4—6.5 cm broad, apex acuminate, base cordate, 3-lobed; central lobe larger, ovate, 6—8 cm long, with margin unequally coarsely dentate, lateral lobes smaller, obliquely flabellate, unequally 2-lobulate, with margin unequally coarsely dentate; surfaces adaxially sparsely strigillose, abaxially appressed-pubescent; venation trinerved, with nerves nearly flat; petioles 4—6.5 cm long, sparsely puberulous. Dichotomous cyme terminal (?), ca. 15 cm broad, three times branched, ca. 12-flowered;




图 1 东川铁线莲 A. 部分枝条；B. 聚散花序；C. 萼片(背面观)；D. 雄蕊；E. 心皮。(孙英宝绘图)

Fig. 1 *Clematis dongchuanensis* A. Part of a branch; B. Cyme; C. Sepal (dorsal view); D. Stamen; E. Carpels. (Drawn by SUN Ying-Bao)

peduncle ca. 7 cm long, subglabrous; bracts 2 opposite, lanceolate-linear, ca. 8 mm long, 1 mm broad, abaxially sparsely pubescent, base decurrent and auriculate;

branches and pedicels glabrous; bracts of branches opposite, linear, 2.5—5.8 mm long, puberulous; pedicels 0.35—6.5 cm long. Flower slightly pendulous: Calyx

broadly campanulate; sepals 4, erect, yellow-green (according to the collector), oblong-ovate, 13—15 mm long, 5—6 mm broad, adaxially and abaxially densely appressed-puberulous, apex obtuse, margin above velutinous. Stamens ca. 50, 11—13.2 mm long; filaments narrow-linear, 9—10 mm long, ca. 0.4 mm broad, densely puberulous, longitudinally 1-nerved; anthers linear, 2.2—2.8 mm long, 0.4 mm broad, abaxially puberulous, apex obtuse. Carpels ca. 40, ca. 9 mm long; ovaries ca. 1 mm long, appressed-pubescent; styles slender, ca. 8 mm long, below densely long sericeous (with hairs 2—4.5 mm long), above densely sericeous-pubescent (with hairs 0.5—1 mm long), apex slightly thickened.

木质藤本。茎长1—2 m(据采集者);小枝细,粗约1 mm,疏被短柔毛。叶对生,为单叶;叶片纸质,心状卵形或五角状卵形,长9—12 cm,宽4—6.5 cm,顶端渐尖,基部心形,3浅裂,中央裂片较长,卵形,长6—8 cm,边缘有不等大的粗牙齿,侧裂片较小,斜扇形,不等微2裂,边缘有不等大的粗牙齿,上面疏被短糙伏毛,下面被贴伏柔毛,三出脉,脉近平;叶柄长4—6.5 cm,疏被短柔毛。二歧聚伞花序顶生(?),宽约15 cm,三回分枝,约有12花;花序梗长约7 cm,近无毛;苞片2,对生,披针状条形,长约8 mm,宽1 mm,下面疏被柔毛,基部下延,呈耳状;分枝和花梗无毛;分枝的苞片对生,条形,长2.5—5.8 mm,被短柔毛;花梗长0.35—6.5 cm。花稍下垂:花萼宽钟状;萼片4,直立,黄绿色(据采集者),长圆状卵形,长13—15 mm,宽5—6 mm,两面密被贴伏

短柔毛,顶端顿,边缘上部被短绒毛。雄蕊约50,长11—13.2 mm;花丝狭条形,长9—10 mm,宽约0.4 mm,密被短柔毛,具1条纵脉;花药条形,长2.2—2.8 mm,宽0.4 mm,背面被短柔毛,顶端钝。心皮约40,长约9 mm;子房约1 mm,被贴伏柔毛;花柱细长,长约8 mm,下部密被长绢毛(毛长2—4.5 mm),上部密被绢状柔毛(毛长0.5—1 mm),顶端稍变粗。

云南(Yunnan):东川(Dongchuan),因民(Yinmin),小山脑(Xiaoshannao),alt. 2290 m,崖边(by cliff),藤本长1—2 m(vine 1—2 m long),花黄绿色(fls. yellow-green),1984-09-11,蓝顺彬(Lan Shun-bin)217(holotype,PE).

本种的叶为单叶,萼片直立,雄蕊花丝和花药均被短柔毛,曲柄铁线莲(*Clematis repens* Finet & Gagnep.)也具上述形态特征,这种情况说明本种在亲缘关系上接近曲柄铁线莲,与后者的区别是本种的叶呈心状卵形,3浅裂,边缘有粗牙齿,花序为具约12朵花的二歧聚伞花序,萼片两面密被短柔毛,雄蕊花药顶端钝,无短尖头。在曲柄铁线莲,其单叶或有时为三出复叶的小叶均呈椭圆形,不分裂,基部宽楔形或圆形,稀浅心形,边缘具少数小牙齿或全缘,聚伞花序腋生,只具1花,萼片外面无毛,雄蕊花药顶端有短尖头(方明渊,1980)。

**致谢** 深切感谢孙英宝先生为本文绘图。

## 参考文献:

Fang MY(方明渊). 1980. *Clematis* sect. *Viorna* [M]//Flora of China(中国植物志). Beijing(北京): Science Press(科学出版社),**28**:88—133

(上接第332页 Continue from page 332)

munity structure in a forest ecosystem in the Southern Mountains of Urumqi(乌鲁木齐南部山区森林生态系统树生地衣群落)[J]. *Chin Bull Bot*(植物学报),**44**(5):578—586

Tumur A(艾尼瓦尔·吐米尔),Abbas A(阿不都拉·阿巴斯). 2006. Floor lichen diversity under different vegetation types in Two-river sources Natural Reserve in Altay Mountains, Xinjiang(新疆阿勒泰山两河源自然保护区地面生地衣的物种多样性)[J]. *Biodivers Sci*(生物多样性),**14**(5):444—450

Tumur A(艾尼瓦尔·吐米尔),Abdulla A(阿地力江·阿不都拉),Abbas A(阿不都拉·阿巴斯). 2011. Distribution pattern of floor lichen communities in Southern mountainous area of Urumqi,Xinjiang,China(乌鲁木齐南部山区地生地衣群落分布

格局)[J]. *Chin Biodivers*(生物多样性),**19**(5):574—580

Wang YL(王玉良),Abdulla A(阿地力江·阿不都拉),Tumur A(艾尼瓦尔·吐米尔),et al. 2012. Diversity and community quantitative characteristics of floor lichen in Yaoluoping National Nature Reserve of Anhui,China(鹞落坪自然保护区地面生地衣多样性及群落数量特征)[J]. *Plant Divers Resour*(植物分类与资源学报),**34**(2):199—207

Yang HX(阳含熙),Lu ZY(卢泽愚). 1981. Methodology of Quantitative Taxonomy in Plant Ecology(植物生态学数量分析方法)[M]. Beijing: Sciences Press:90—252

Zedda L,Rambold G. 2009. Diversity and ecology of soil lichens in the Knersvlakte(South Africa)[J]. *Bryologist*,**112**:19—29